

**ANALYTICAL METHOD VALIDATION COMMITTEE FOR NON
PHARMACOPOEIAL PRODUCT
DEPARTMENT OF DRUG ADMINISTRATION
National Medicines Laboratory**

Rifaximin tablet

Analytical Profile No.: RIF 074/075/AP 026

Rifaximin tablets contain not less than 90 per cent and not more than 110 per cent of the stated amount of Rifaximin.

1. Identification:

In the assay, the principle peak in the chromatogram obtained with the sample solution should correspond to the peak in the chromatogram obtained with the reference standard solution of Rifaximin.

2. Dissolution Test: Determine by liquid chromatography

2.1 Dissolution Parameters:

Apparatus: Paddle

Medium: 1000 ml of 0.1 M sodium phosphate buffer pH 7.4 containing 0.8 per cent sodium lauryl sulphate

Speed and time: 75 rpm and 60 minutes

Withdraw the suitable volume of the medium and filter.

2.2 Test solution:

Dilute 5 ml of the filtrate to 25 ml with mobile phase and filter through 0.22 micron filter paper.

2.3 Reference solution:

Weigh accurately about 27.5 mg of working standard of Rifaximin and transfer into 50 ml volumetric flask. Dissolve with mobile phase and make up the volume to 100 ml with mobile phase. Pipette 5 ml of this solution and transfer into 25 ml volumetric flask, add 5 ml of dissolution medium and make up the volume to 25 ml with mobile phase. Filter through 0.22 micron filter paper.

2.4 Chromatographic system:

Use the chromatographic system as described under assay

**ANALYTICAL METHOD VALIDATION COMMITTEE FOR NON
PHARMACOPOEIAL PRODUCT
DEPARTMENT OF DRUG ADMINISTRATION
National Medicines Laboratory**

2.5 Procedure:

Inject 20 µl of reference and test solution five/six times and obtain the respective chromatogram. Measure the peak responses. Calculate the % release.

2.6 Limit:

D. not less than 80 per cent of the stated amount

3. Assay: Determine by liquid chromatography

3.1 Buffer:

0.025 M sodium dihydrogen phosphate in water, adjust pH to 3.0 with orthophosphoric acid.

3.2 Test Solution:

Weigh individually 20 tablets and crush the tablet to fine powder. Weigh accurately the powder equivalent to 50 mg of rifaximin and transfer into 50 ml volumetric flask. Add about 35 ml of diluent, sonicate for about 10 minutes and cool the solution to room temperature and make up the volume to 50 ml with diluents. Centrifuge the solution. Dilute 5 ml of the resulting solution to 50 ml with diluent. Filter the solution with 0.22 micron membrane filter paper.

3.3 Reference solution:

Weigh accurately about 50 mg of working standard of Rifaximin and transfer into 50 ml volumetric flask. Dissolve in the mobile phase and make up the volume to 50 ml with mobile phase. Dilute 5 ml of the resulting solution to 50 ml with mobile phase. Filter through 0.22 micron membrane filter paper.

3.4 Chromatographic system:

Column: Phenyl column, 25 cm x 4.6 mm

Wavelength: 300nm

Flow rate: 1.0 ml per minute

Injection volume: 20 µl

Column temperature: 35 °C

Detector: UV

Mobile phase:

A mixture of 45 volumes of buffer solution (0.025 M sodium dihydrogen phosphate in water, adjust pH to 3.0 with orthophosphoric acid.) and 55 volumes of acetonitrile.

**ANALYTICAL METHOD VALIDATION COMMITTEE FOR NON
PHARMACOPOEIAL PRODUCT
DEPARTMENT OF DRUG ADMINISTRATION
National Medicines Laboratory**

3.5 Procedure:

Inject 20 µl of standard preparation five/six times. The test is not valid unless the column efficiency determined from the major peak is not less than 2000 theoretical plates, the tailing factor is not more than 2.0 and the relative standard deviation of replicate injections is not more than 2.0 %.

Inject 20 µl of standard and sample solution separately and obtain the respective chromatogram. Measure the peak responses. Calculate the content of rifaximin per tablet.

4. Other tests: As per pharmacopoeial requirements.